

# The Global Land Surface Model ORCHIDEE

(ORganizing Carbon and Hydrology In Dynamic Ecosystems Environment)



Presented by Nicolas Vuichard  
LSCE/IPSL


# Outline

---

- ★ A brief history of ORCHIDEE & motivations
- ★ Formalism
- ★ Main processes
- ★ Configurations & Inputs requirements


# A brief history

Model



(Laval et al., 1981)      (Ducoudré et al., 1993)      (Viovy et al., 1997)      (Polcher et al., 1998)      (Krinner et al., 2005)

Project / Users



# Objective

---

- ★ Simulate Energy, Water and Carbon fluxes at the land surface/ atmosphere interface.
  - To be used for being the 'land surface' component of a Earth system model (IPSL-CM5).
    - Global => to represent the main vegetation cover.
    - Regional => to study feedback processes.
  - For past, present and future climates
    - Module of vegetation dynamic
    - Process-based modeling
- ★ Conservation of mass and energy is a guiding principle for ORCHIDEE.


# Outline

---

- ★ A brief history of ORCHIDEE & motivations
- ★ **Formalism**
- ★ Main processes
- ★ Configurations & Inputs requirements

# Surface variability representation ?

★ A mosaic of vegetation and soil moisture



# Surface variability representation ?

---

★ In each grid cell, we account for:

★ Bare soil

• *veget\_max(1)*

★ Vegetated lands

• *veget\_max(2:nvm)*

★ Other lands (so far, only the continental ice)


• *frac\_nobio*

$$\sum_{i=1}^{nvm} (\text{veget\_max}_i) + \text{frac\_nobio} = 1$$

★ One soil type per grid cell but different soil moisture profiles.

# Vegetated lands

- ★ Concept of 'Plant Functional Types' (PFT)
- ★ Defined according to systematic, physiological, phenological, climatic conditions






# Plant Functional Types

- ★ A same set of equations governs PFT
- ★ But parameter values differ among PFT's

PFT	$V_{cmax,opt}$	$T_{opt}$	$\lambda_{max}$	$Z_{root}$	$\alpha_{leaf}$	$h$	$A_c$	$T_s$	$H_s$
TrBE	50	37	10	1.25	0.12	25	910	-	0.3
TrBR	60	37	10	1.25	0.14	25	180	-	0.3
TeNE	37.5	27	5	1.	0.14	15	910	-	-
TeBE	37.5	32	5	1.25	0.14	15	730	-	-
TeBS	37.5	28	5	1.25	0.14	15	180	12.5	-
BoNE	37.5	25	4.5	1.	0.14	10	910	-	-
BoBS	37.5	25	4.5	1.	0.14	10	180	5	-
BoNS	35	25	4	1.25	0.14	10	180	7	-
NC3	70	$27.5 + 0.25T_l$	2.5	0.25	0.20	0.2	120	4	0.2
NC4	70	36	2.5	0.25	0.20	0.2	120	5	0.2
AC3	90	$27.5 + 0.25T_l$	6	0.25	0.18	0.4	150	10	0.2
AC4	90	36	3	0.25	0.18	0.4	120	10	0.2

# Response to environmental conditions



# Concept of externalization

---


- ★ By default 13 PFT's (named Metaclass) with pre-defined parameters setting
- ★ Most of the parameters can be modified by the user (see <http://forge.ipsl.jussieu.fr/orchidee/wiki/Documentation/OrchideeParameters> or orchidee.default file in the config/PARAM directory)
- ★ The number of PFT's can be extended
  - ★ By setting the NVM parameter and PFT\_TO\_MTC (correspondance array linking a PFT to MTC)

# Outline

---

- ★ A brief history of ORCHIDEE & motivations
- ★ Formalism
- ★ **Main processes**
- ★ Configurations & Inputs requirements

# Main processes



# Energy budget & Resistance terms

## *enerbil module*

- ★ Calculation of :
  - ★ Sensible heat flux
  - ★ Latent heat flux
    - Transpiration
    - Evaporation of bare soil and leaf water
    - Sublimation
  - ★ Net radiation
  - ★ Soil and surface temp.
  - ★ GPP calculation
- ★ *One calculation per grid cell*
- ★ *No vertical discretization*

## *diffuco module*


- ★ vbeta1 : sublimation
- ★ vbeta2 : interception loss
- ★ vbeta3 : transpiration
- ★ vbeta4 : bare soil evaporation
- ★ vbeta5 : flood plains

# C assimilation/stomatal conductance


*diffuco module*  
*diffuco\_trans\_co2 routine*

- ★ A and  $G_s$  are calculated at each LAI level:
- ★ Beer-Lambert decrease of light in the canopy
- ★ Exponential decrease of  $V_{max}$  (but limited to 30%) to mimic nitrogen decrease
- ★ The others parameters (e.g  $CO_2$ , rel hum..) are held constants.

From the leaf to canopy




Light profile



# Vertical discretization in the soils

ORCHIDEE used to have different vertical discretizations for moisture and temperature. The physics require different numerical choices !

This was not tenable any more with soil freezing processes, permafrost and complex snow schemes.



★ The higher vertical resolution does not deteriorate the thermal diffusion (CFL criteria !)

★ Users are now provided with a set of parameters to configure the soils.

- zmax\_t (DEPTH\_MAX\_T = 10 )
- zmax\_h (DEPTH\_MAX\_H = 2 )
- depth\_topthickness ( ~1 mm)
- refinebottom
- ratio\_geom\_below




# Soil temperatures

---

## *thermosoil module*

- ★ Calculates the soil temperatures by solving the heat diffusion equation within the soil
  - the soil is divided into several layers, reaching at least 10m down within the soil. The user can adapt the model to the application.
  - Thickness follows a geometric series.
  - No more interpolation with soil water layers


- ★ Physically-based description of soil water fluxes using Richards equation : 2m soil discretized in at least 11-layers.
- ★ Hydraulic properties based on van Genuchten-Mualem formulation
- ★ Related parameter based on texture (fine, medium, coarse)
- ★ Surface runoff =  $P - E_{sol} - \text{Infiltration}$
- ★ Free drainage at the bottom




# Routing / Irrigation

*routing module*

- ★ Routing parametrization to calculate water discharge to river



*From Guimberteau (thesis, 2010)*



*The 50 major river basins on the LMD-GCM grid*

# Biomass and soil pools

---

- ★ 8 pools of living biomass

- ★ Leaves, fine roots, above and below sapwood, above and below heartwood, 'fruits' and 'reserves'

- ★ 4 pools of litter

- ★ Above/below, Structural & Metabolic

- ★ 3 pools of soil


- ★ Active, Slow and Passive

# Allocation of assimilates


*stomate\_alloc module*

- ★ Principle of resource optimisation
- ★ Allocation to leaves, branches, roots as a function of resources : water (H), nitrogen (N) and light (L)


Leaf = 1 - Branch - Root



$$\text{Branch} = \frac{3 T_0 \min(H,N)}{2 L + \min(H,N)}$$



$$\text{Root} = \frac{3 R_0 L}{L + 2 \min(H,N)}$$



# Phenology

---

*stomate\_phenology module*

- ★ Bud-burst model (Botta et al. 2000)
  - ★ Defined for each PFT based on Growing degree days, Number of chilling days, soil water, ...
  - ★ Calibrated at global scale from bud-burst estimated by satellite
- ★ Senescence
  - ★ Function of leaf age and environmental conditions *stomate\_turnover module*
  - ★ For trees, a senescence stage is considered until all leaves fall (while for grass senescence it is a continuous process)

# Respirations

## ★ Autotrophic respiration

### ★ Maintenance

- linear response to temperature (Ruimy et al.)
- potential adaptation to long term temperature

### ★ Growth

- a fixed part of assimilates


*stomate\_npp module*

## ★ Heterotrophic respiration

- Century-like model

*stomate\_litter & stomate\_soilcarbon modules*

*stomate\_resp module*



# Land cover change

---

*stomate\_lcchange module*

- ★ Vegetation map can vary from one year to another
  - ★ For decreasing PFT
    - A part of biomass is exported and goes to 3 decomposition pools (1 year, 10 year 100 years), the rest goes to litter of increasing PFTs.
    - Soil carbon of decreasing PFT is diluted to soil carbon of increasing PFT's
  - ★ When a PFT disappears => reset to its initial state
  - ★ When a PFT appears => growth from seed



# Vegetation dynamic

---

*lpj\_kill, lpj\_pftinout, lpj\_constraints modules*


- ★ Taken from LPJ model
- ★ All PFT's are able to growth in each grid cell
  - Climate constraints define regeneration and adaptation of PFT's
  - Light competition when canopy closure (PFT with NPPmax dominate)
  - Trees always dominate grasses

# Outline


---

- ★ A brief history of ORCHIDEE & motivations
- ★ Formalism
- ★ Main processes
- ★ **Configurations & Inputs requirements**

# Tasks performed by ORCHIDEE for the IPSL ESM



# Infrastructure surrounding ORCHIDEE




- Some numbers to ORCHIDEE's code :
- 83 FORTRAN files
  - 61000 lines of code
  - 43000 lines of comments

# Structure of the code

---

- ★ Use of a modular structure
  - ★ All the variables are dynamics (allocatable)
  - ★ For each module:
    - A main entry point : <module>\_main
    - An initialisation procedure : <module>\_initialize
    - An end procedure : <module>\_finalize
    - An procedure to clear memory : <module>\_clear
  - ★ All the variables are transmitted by subroutine parameters.
  - ★ Prognostic variables are local to the modules.

# Subroutine Call graph





SECHIBA(30min)  
 STOMATE(30min)  
 STOMATE(1 day)

30<sup>PJ</sup>

# Atmospheric Interface

★ Meteorological forcing (from monthly to half-hourly)



# Forcing files

---

## ★ Meteorological data

- ★ One often uses reanalysis or in-situ data with different time resolution (3h, 6h, 1/2 hour, ...)
- ★ The spatial resolution of the simulation is driven by the resolution of the meteo forcing file.
- ★ The time step of a simulation is defined by the parameter TIME\_STEP (30 min by default).
- ★ The meteorological data often needs to be interpolated in time to the ORCHIDEE time step.



# Ancillary data

---

- ★ Ancillary data needed will depend on the configuration chosen.
- ★ All variables will be interpolated to the grid of ORCHIDEE.
- ★ Some exemples :
  - PFT map and land use
  - soil texture
  - Background albedo
  - Topographic slopes
  - River graphs

# Conclusions

---

- ★ ORCHIDEE is a complex system !
- ★ But you have the chance to use a system which was developed at IPSL and by people who are still present.
- ★ The model has too many options and you will get lost!
- ★ Do not hesitate to ask the original developers if you have problems.
- ★ Enjoy the training !